

According to Rudolph and other's study, while oxygen in the water was increased in winter due to the dominant of the north-west wind, in the summer, it was reduced by the water input from outside the San Vicente Bay<sup>17</sup>. According to Karthik and other's study, high dissolved oxygen and low salinity at seawater of Port Blair, South Andaman Island during September and November was caused by rainfall – induced freshwater input and flooding from land<sup>18</sup>. Both abundance and diversity of finfish, lobster and squid was significantly decreased when dissolved oxygen was less than 2 mg/L in bottom of western Long Island Sound<sup>19</sup>.

In this study, temperature, pH, and concentrations of arsenic (As), chromium (Cr), nickel (Ni), lead (Pb) and zinc (Zn) between 19 October 2016 and 4 June 2017 and temperature, pH, dissolved oxygen from 18 November 2018 to 31 January 2019 were investigated in seawater samples collected from Kucusu, Fenerbace and Suadiye beaches in Istanbul on weekly. The objective of the study was to assess the possible effects of seawater on public health.

### Experimental

100 ml-polyethylene bottles were soaked in a mixture of nitric acid and deionized water (Fig. 1a) for at least 24 h, were rinsed three times with deionized water and, then, were dried. Seawater samples were collected on a weekly basis at 50 cm below the sea-surface in these polyethylene bottles, which were attached to a 3–4 m long telescopic rod<sup>20</sup>. Before filling the bottles, they were being rinsed three times with seawater, and then they were filled completely. The temperature, dissolved oxygen and pH of seawater were determined onsite using a multiparameter (YSI) (Fig. 1b). Then, the samples were delivered refrigerated to the laboratory of Marmara University for analysis. In the laboratory, each

sample was filtered through a 0.45  $\mu\text{m}$  sieve syringe filter (Sartorius) using a sterile syringe. The samples were diluted using a dilution factor of 100 (Fig. 1c). To protect the metals and prevent sedimentation, the pH of the samples was adjusted to below 2 using a 2% nitric acid ( $\text{HNO}_3$ ). Then, the samples were stored in a 4°C refrigerator until its analysis using ICP-MS (Agilent 7500A) (Fig. 1d) to determine the concentration of heavy metals.

### Results and discussion


The temperature and pH values measured between 19 October 2016 and 26 March 2017 ranged from 11.8 to 19.2°C and from 7.1 to 9, respectively (Fig. 2a). The maximum pH value was the one measured on 26 October 2016 in the Suadiye beach and it was below the threshold values for Turkey<sup>21,22</sup> (6–9) and EPA<sup>23</sup> (5–9 for human health, and 6.5–8.5 for aquatic life). In addition, higher values of pH were observed in the direction of the south-east wind. On the other hand, due to snowing, the lowest sea temperature was observed in the middle of January.

In Fenerbahce, pH ranged from 7.55 to 7.89, with a higher range of 8.01 to 8.51 in Kucusu, with a lower range of 8.01 to 8.32 in Suadiye from 18 November 2018 to 31 January 2019 (Fig. 2b). The weekly water temperatures from the three sites were from 6.3 to 15.4°C and the weekly dissolved oxygen was in the range of 80 to 95.4% from 18 November 2018 to 31 January 2019 (Fig. 3a). These dissolved oxygen values did not exceed the Turkish threshold values (in the range of 80 to 120%).


The highest pH was in Kucusu as 8.51 on 25 November 2018, and the highest temperature was in Kucusu as 15.4°C on 18 November 2018, and the highest dissolved oxygen was in Suadiye as 95.4% in 25 November 2018. While sea-



**Fig. 1.** (a) Elga Purelab Flex Deionized water, (b) YSI Professional plus multiparameter meter, (c) Sampling bottles and vials, (d) Autosampler (CETAC ASX-51) and ICP-MS (Agilent 7500a).



**Fig. 2.** Weekly temperature, pH values (a) between 19 October 2016 and 26 March 2017 and (b) between 18 November 2018 and 31 January 2019.



**Fig. 3.** (a) Weekly dissolved oxygen values and (b) Cr concentrations between 18 November 2018 and 31 January 2019.

water temperatures of Kucuksu, Fenerbahce, and Suadiye beaches were low in January 2019, they were high in November 2018.

Strength of association for Pearson correlation was investigated<sup>24</sup>. For Kucuksu and Fenerbahce ( $-0.67$ ) and for Kucuksu and Suadiye ( $-0.58$ ), pH had strong negative correlations but pH had strong positive correlation ( $0.70$ ) between Fenerbahce and Suadiye. Therefore, when the pH value increased for Fenerbahce and Suadiye, pH value decreased for Kucuksu. Temperature correlations were greater than  $0.87$  and were strong for all beaches like dissolved oxygen correlations. Furthermore, for all beaches, seawater temperatures decreased until middle of January 2019 and later temperatures increased. The dissolved oxygen values synchronously decreased for all beaches.

pH and temperature correlation was  $0.28$  for Kucuksu,  $0.10$  for Fenerbahce, and  $0.12$  for Suadiye and these correlations were weak for all beaches. For Suadiye, pH and dissolved oxygen had strong positive correlation ( $0.68$ ). There was a positive correlation between temperature and dissolved oxygen in the range from  $0.40$  to  $0.78$ . This means that the dissolved oxygen was affected by water temperature like

Hamzah and others study<sup>15</sup>.

pH range in our study is wider than study in Malaysia ( $7.59$ – $7.8$ ) but pH ranges in both studies are suitable for human health and aquatic life because Turkish pH standard is  $6$ – $9$  and Malaysia pH standard is  $6.5$ – $8.5$ . Minimum temperature in our study is less than studies in Malaysia ( $18.9^{\circ}\text{C}$ ), Chile ( $10.59^{\circ}\text{C}$ ), and India ( $25^{\circ}\text{C}$ ).

Minimum dissolved oxygen in our study ( $8.8$  mg/L) is higher than minimum values in Malaysia ( $3.7$  mg/L), and India ( $3.2$  mg/L). So this vital parameter in marine life positively affects the Sea of Marmara in terms of aquatic creature's diversity and number.

The concentrations of As, Cr, Ni, Pb and Zn (Fig. 3b, 4a,b) were found to be lower than the threshold values<sup>22</sup> ( $0.1$  mg/L for every element). In addition, the values were found to be lower in windy weather. For example, on October 26 and 12 December 2016, all element concentrations were found to be relatively low compared to the values measured during the previous week.

An assessment was conducted to estimate the correlations between the concentrations of each element with temperature. A negative correlation ( $-0.4$ ) was found between arsenic and temperature. This indicates that the arsenic concentration increases as the temperature decreases. On the other hand, the positive correlation ( $0.5$ ) between temperature and chromium shows that its concentration increases with temperature. The Pb concentrations in seawater of Fenerbahce beach on 16 November 2016 and in Kucuksu beach on 1 April 2017 were high; however, they did not exceed the threshold value.

Zn concentrations were also lower than the threshold value, although they were relatively high at the end of Janu-




Fig. 4. Weekly heavy metal concentrations: (a) Ni and As and (b) Zn and Pb between 18 November 2018 and 31 January 2019.

to be lower than that of the studies held in Mexico<sup>9</sup> and Turkey<sup>25</sup> as well as the maximum allowable limit in Mexico. Similarly, the maximum Pb value (0.1 ppm) was found to be lower than the maximum allowable limits and the maximum values measured in studies carried out in Mexico<sup>9</sup>, Indonesia<sup>7</sup> and Turkey<sup>25</sup>. Finally, the maximum zinc value was found to be close to the values measured in the studies carried out in Mexico and Indonesia and below the maximum allowable

Table 1. Comparing with references about heavy metals

Element	Concentration (ppm)				Country for limit value	Coast	Year
	Sampling point A	Sampling point B	Sampling point C	Limit			
Zn <sup>7</sup>	0.155	0.126	0.109	0.01	Indonesia	Mamboro District of North Palu	2015
Zn (max) <sup>9</sup>	0.167	0.153	0.175	20	Mexica	Mexican Pasific Ocean coastline	2013
Zn (max) <sup>25</sup>	5.45			0.1	Turkey	Black Sea	2010
Pb <sup>7</sup>	0.35	0.391	0.433	0.025	Indonesia	Mamboro District of North Palu	2015
Pb (max) <sup>25</sup>	0.808			0.1	Turkey	Black Sea	2010
As (max) <sup>9</sup>	0.001	0.001	0.001	0.1	Mexica	Mexican Pasific Ocean coastline	2013
Cd (max) <sup>9</sup>	0.093	0.093	0.091	0.1	Mexica	Mexican Pasific Ocean coastline	2013
Cd (max) <sup>25</sup>	0.169			0.01	Turkey	Black Sea	2010
Cu (max) <sup>9</sup>	0.044	0.044	0.044	1	Mexica	Mexican Pasific Ocean coastline	2013
Cu (max) <sup>25</sup>	0.775			0.01	Turkey	Black Sea	2010
Cr (max) <sup>9</sup>	0.146	0.097	0.097	0.5	Mexica	Mexican Pasific Ocean coastline	2013
Cr (max) <sup>25</sup>	0.582			0.1	Turkey	Black Sea	2010
Ni (max) <sup>9</sup>	0.325	0.399	0.381	2	Mexica	Mexican Pasific Ocean coastline	2013
Ni (max) <sup>25</sup>	0.833			0.1	Turkey	Black Sea	2010
Pb (max) <sup>9</sup>	0.373	0.356	0.349	0.5	Mexica	Mexican Pasific Ocean coastline	2013

ary and the beginning of February; especially, during the first week of February, because of the fuel oil leakage at the Gulf of İzmit in the Sea of Marmara that occurred on 12 January 2017<sup>13</sup>.

The maximum arsenic value (0.03 ppm) was compared to the value measured in a study held in Mexico<sup>9</sup> (0.001 ppm) and was found to be higher (Table 1). However, it was still lower than the maximum allowable limit in Mexico. The maximum Cr value (0.102 ppm) measured in the present study was found to be close to that measured in the same Mexican study, which is lower than the maximum allowable limit in Mexico<sup>12</sup>, and it was also lower than the Cr value (0.58 ppm) measured in a similar study in Turkey<sup>25</sup>. On the other hand, the maximum nickel value (0.02 ppm) in this study was found

limits in Mexico, Indonesia and Turkey.

Briefly, compared to previous studies carried out in Mexico and the Black Sea, Turkey, the concentrations of Cr, Ni, Pb and Zn in the present study were lower. On the other hand, Pb and Zn concentrations in a study carried out in Indonesia were higher than the values measured in the present study. Finally, the results showed that the concentrations of all the elements measured in this study were below the maximum allowable limits in Mexico and Indonesia.

### Conclusions

The results of the analysis showed that the level of pH and dissolved oxygen and the heavy metal (As, Cr, Ni, Pb and Zn) concentrations in the Sea of Marmara did not repre-

sent a serious threat to the aquatic life and human health, because they were all below the threshold values. It was observed that the concentrations of all elements decreased in October 26th and December 20th, when the wind was excessive.

The lowest sea temperature between 19 October 2016 and 26 March 2017 was observed in the middle of January due to snow. In addition, the temperature was found to be positively correlated with the arsenic concentration and was negatively correlated with the chromium concentration. The maximum Pb concentrations were observed at the Fenerbace beach on 16 November 2016 and at the Kukuksu beach on 1 April 2017. There was an increase in the zinc concentrations in the Sea of Marmara, where the fuel leak occurred in the Gulf of İzmit on 12 January 2017.

At all three beaches from 18 November 2018 to 31 January 2019, seawater was warmer on the January, 2019 while the seawater was colder in November, 2018 than other months. The range of seawater temperatures recorded in this study was the normal for humid subtropical climate. Correlation values indicated that the dissolved oxygen was affected by water temperature.

The concentrations of all the tested elements except As were found to be low compared to studies carried out in Mexico and the Black Sea, Turkey. In addition, the concentrations of all tested elements were found to be below the Mexican limits. Furthermore, the Pb and Zn concentrations in the current study were lower than those measured in the study conducted in Indonesia.

These results show that, to enable immediate actions for the mitigation of accidents and natural disasters, beaches should be monitored periodically.

### Acknowledgements

This work was supported by Research Fund of the Marmara University. Project Number: FEN-A-110718-0394, and FEN-C-110718-0405.

### References

1. A. Yanez-Arancibia and P. Sanchez-Gil, in: "Ecología de los recursos demersales marinos (AGT Editor)", DF, 1988, 189.
2. D. J. Jefferies and P. Freestone, *Journal of Otter Trust*, 1984, **18**, 17.
3. R. K. Singh, S. L. Chavan and P. H. Sapkale, *Environmental Monitoring and Assessment*, 2007, **129**, 471.
4. A. G. Cardoso, G. R. Boaventura, E. V. S. Filho and J. A. Brod, *Journal of the Brazilian Chemical Society*, 2001, **12**, 767.
5. H. A. Naser, *Marine Pollution Bulletin*, 2013, **72**, 6.
6. F. Aydinol, I. Turkdogan, G. Kanat and H. Bayhan, *Environmental Monitoring and Assessment*, 2012, **184**, 149.
7. T. Rahmadani, S. M. Sabang and I. Said, *Jurnal Akademika Kimia*, 2015, **4**, 197.
8. E. Bozkurt, O. Eliri and M. Kesiktas, *Journal of Recreation and Tourism Research*, 2014, **1**, 39.
9. V. Perez-Moreno, M. A. Ramos-López, C. E. Zavala-Gomez and M. A. Rico Rodriguez, *Interciencia*, 2016, **41**, 419.
10. I. I. Ozyigit, O. L. Uyanik, N. R. Sahin, I. E. Yalcin and G. Demir, *Polish Journal of Environmental Studies*, 2017, **26**, 773.
11. F. Solomon, *Environment and Communities*, 2009, **15**, 25.
12. Nytimes, 2018. <https://www.nytimes.com/aponline/2017/07/18/world/europe/ap-eu-turkey-heavy-rainfall.html>.
13. World Maritime News, 2018. <http://worldmaritimeneews.com/archives/210453/fuel-oil-spills-at-izmit-bay-traffic-halted/>.
14. WHO (World Health Organization), in Guidelines for safe recreational water environments, WHO, 2003, Executive summary, 28.
15. A. Hamzah, S. H. Kipli, S. R. Ismail, R. Una and S. Sarmani, *Sains Malaysiana*, 2011, **40**, 93.
16. State of Israel, in Environmental quality standards for the Mediterranean Sea in Israel, State of Israel, 2002, 22.
17. A. Rudolph, R. Ahumada and C. Pérez, *Environmental Monitoring and Assessment*, 2002, **78**, 89.
18. R. Karthik, M. Arun Kumar, S. Sai Elangovan, R. Siva Sankar and G. Padmavati, *J. Mar. Biol. Oceanogr.*, 2012, **1**, 1.
19. P. Howell and D. Simpson, *Estuaries*, 1994, **17**, 394.
20. G. E. Batley, in: "Trace Element Speciation Analytical Methods and Problems", ed. G. E. Batley, CRC Press, 1989, 3.
21. State of Turkey, in: "Turkey Recreational Standards", State of Turkey, 2006, Annexes 1, 1.
22. State of Turkey, in Turkish Water Pollution Control Regulation, State of Turkey, 2004, Annexes, Table 4.
23. U.S. EPA, in Quality criteria for water, U.S. EPA, 1986, Appendix A, 233.
24. E. Bozkurt Kopuz, A. Pursa, B. Yilmaz, E. Ozdemir and D. Ozturk, in: "Current Trends in Science and Landscape Management", eds. R. Efe, M. Zencirkiran, J. A. Wendt, Z. Tumsavas, H. Unal and B. Borisova, St. Kliment Ohridski University Press, 2017, **33**, 417.
25. M. Boran and I. Altinok, *Turkish Journal of Fisheries and Aquatic Sciences*, 2010, **10**, 565.

